

**WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018**

Drug classes not included on this list are not managed through a Preferred Drug List (PDL).
HOWEVER, THIS EXCLUSION IS NOT A GUARANTEE OF PAYMENT OR COVERAGE. Dosage limits and other requirements may apply.
Drugs new to market are non-preferred until a clinical review has been completed. PA criteria will apply to both the pediatric population,
as well as the adult population for those plans where PA/PDL limits are allowed.
Unless otherwise noted on the PDL, generic substitution is mandatory.
Yellow highlighted items below indicate new changes to the PDL. Red font indicates quantity/dosage limits apply. *Indicates BRAND is Preferred. May Use DAW 5.
Contact the Change Healthcare PA Helpdesk @ 877-207-1126 for prior authorization if client has primary insurance that will not cover the brand name medication.

Please refer to the Additional Therapeutic Criteria Chart, **Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.**

Therapeutic Class	Preferred Agents	Preferred Agents Requiring Clinical Criteria	Clinical Criteria	Non-Preferred Agents Generic Mandatory Policy Applies <small>(THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS)</small>
ADDICTION	BUPRENORPHINE COMBINATIONS	SUBOXONE FILM	Client must have a diagnosis of opioid dependence or abuse. This is not to be used for the treatment of chronic pain. Prescriber must have a XDEA number. Prior authorization will be required before any narcotic, benzodiazepine, or carisoprodol prescription will be allowed between fills. Prior authorization will be required before any short-acting stimulant prescription from any doctor other than the prescriber of buprenorphine or Suboxone, will be allowed between fills. Oral buprenorphine will be approved for clients that are pregnant or nursing or with a documented allergy to naloxone. Please submit PA requests on the "Oral Buprenorphine/Naloxone or Oral Buprenorphine" PA form available at www.wymedicaid.org . Dosage limits apply During first two years of treatment: 16mg After two years of treatment: 8mg	BUNAVAIL buprenorphine (oral) buprenorphine/naloxone tablets (use preferred) ZUBSOLV
	NALTREXONE	naltrexone VIVITROL	Client must have a diagnosis of alcohol or opioid dependence. Prior authorization will be required before any narcotic, carisoprodol, or benzodiazepine prescription will be allowed between fills. Prior authorization will be required before a short-acting stimulant prescription from any doctor other than the prescriber of naltrexone or Vivitrol will be allowed between fills.	
ALLERGY / ASTHMA	ANTI-HISTAMINES, MINIMALLY SEDATING	cetirizine fexofenadine loratadine	Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	desloratadine CLARINEX RDT/SYRUP levocetirizine
	ANTI-HISTAMINE/DECONGESTANT COMBINATIONS	cetirizine/pseudoephedrine fexofenadine/pseudoephedrine loratadine/pseudoephedrine	Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	CLARINEX-D
	ANTICHOLINERGIC BRONCHODILATORS	ipratropium SPIRIVA HANDIHALER	Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Spiriva 5 day STARTER package will be allowed one (1) time per recipient.	ATROVENT HFA INCRUSE ELLIPTA SEEBRI SPIRIVA RESPIMAT (use preferred agent) TUDORZA
	INHALED COMBINATION AGENTS	ADVAIR DISK/HFA SYMBICORT	Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. **Will also require the diagnosis of COPD. ***Will also require the diagnosis of COPD or uncontrolled asthma. Advair 7 and 14-day STARTER package will be allowed one (1) time per recipient.	ANORO ELLIPTA** BEVESPI BREQ ELLIPTA*** COMBIVENT DULERA fluticasone/salmeterol 232,113,55-14mcg STIOLTO TRELLEGY UTIBRON
	LEUKOTRIENE MODIFIERS	montelukast	Trial and failure of preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent.	zafirlukast ZYFLO
	LONG ACTING BRONCHODILATORS	BROVANA FORADIL SEREVENT	Trial and failure of two (2) preferred agents greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent.	PERFORMIST STRIVERDI
	NASAL ANTIHISTAMINES	ASTELIN* azelastine 0.1%	Trial and failure of preferred agent greater than or equal to 90 days in the last 12 months will be required before approval can be given for a non-preferred agent.	azelastine 0.15% AZENASE (use separate agents) DYMISTA (use separate agents) olopatadine 0.6%
	NASAL STEROIDS	BECONASE AQ flunisolide fluticasone NASONEX*	Trial and failure of two (2) preferred agents greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Budesonide will be approved for pregnancy.	AZENASE (use separate agents) budesonide DYMISTA (use separate agents) mometasone (BRAND IS PREFERRED) OMNARIS QNASL TICANASE (use separate agents) triamcinolone VERAMYST ZETONNA
	SHORT ACTING BRONCHODILATORS - INHALERS	PROAIR HFA PROVENTIL HFA VENTOLIN HFA	Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Minimum day supply of at 16 days is required	PROAIR RESPICLICK XOPENEX HFA*

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria.				
THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small>
ALLERGY / ASTHMA continued	SHORT ACTING BRONCHODILATORS - NEBULIZERS		Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent.	
	albuterol neb levalbuterol neb			
	STEROID INHALANTS		Trial and failure of two (2) preferred agents greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Alvesco will be approved for a history of oral thrush with steroid inhalants.	AEROBID/AEROBID-M AEROSPAN ALVESCO ARMONAIR ARNUNITY ASMANEX budesonide susp 0.25mg, 0.5mg, and 1mg (BRAND IS PREFERRED) QVAR
EPINEPHRINE				ADRENACLICK (use preferred agent) AUVI-Q (use preferred agent) EPI-PEN (use preferred agent)
ARTHRITIS	IMMUNOMODULATORS		Client must have diagnosis of AS prior to approval of a step 1 agent (Enbrel or Humira). To receive Cosentyx, the client must have a diagnosis of AS and a 56-day trial and failure of Humira. To receive a non-preferred agent, client must have a diagnosis of AS and a 56-day trial and failure of both preferred agents	CIMZIA REMICADE SIMPONI
	ANKYLOSING SPONDYLITIS (AS)			
	STEP 1 AGENTS			
		ENBREL HUMIRA		
	STEP 2 AGENTS		Quantity Limits apply for all diagnoses: Enbrel 25mg - limited to 10 per month Enbrel 50mg - limited to 5 per month Humira 20mg - limited to 10 per month Humira 40mg - limited to 5 per month	
		COSENTYX		
	JUVENILE IDIOPATHIC ARTHRITIS (JIA)		Client must have diagnosis of JIA prior to approval of a preferred agent. To receive a non-preferred agent, client must have a diagnosis of JIA and a 56-day trial and failure of both preferred agents.	ACTEMRA ORENCIA
		ENBREL HUMIRA		
	PSORIATIC ARTHRITIS (PA)		Client must have diagnosis of AS prior to approval of a step 1 agent (Enbrel or Humira). To receive Cosentyx, the client must have a diagnosis of AS and a 56-day trial and failure of Humira. To receive a non-preferred agent, client must have a diagnosis of AS and a 56-day trial and failure of both preferred agents	CIMZIA OTEZLA REMICADE SIMPONI
	STEP 1 AGENTS			
	ENBREL HUMIRA			
STEP 2 AGENTS				
	COSENTYX			
RHEUMATOID ARTHRITIS (RA)		Client must have diagnosis of RA and a 56-day trial and failure of methotrexate prior to approval of a preferred agent. To receive a non-preferred agent, client must have a diagnosis of RA and a 56-day trial and failure of both preferred agents.	ACTEMRA CIMZIA KEVZARA KINERET ORENCIA REMICADE RITUXAN SIMPONI XELJANZ/XR	
	ENBREL HUMIRA			
CONVULSIONS	DIAZEPAM RECTAL GEL			diazepam gel (BRAND IS PREFERRED)
	DIASTAT*			
	ORAL ANTICONSULSANTS		Limited to FDA approved indications	
		APTOM FYCOMPA VIMPAT		
CROHN'S	IMMUNOMODULATORS		Client must have diagnosis of Crohn's prior to approval of the preferred agent. To receive a non-preferred agent, client must have a diagnosis of Crohn's and a 56-day trial and failure of the preferred agent.	CIMZIA REMICADE STELARA TYSABRI (additional criteria applies)
		HUMIRA		
DERMATOLOGY	IMPETIGO ANTIBIOTICS		Trial and failure of ALL preferred agents greater than or equal to 7 days in the past 90 days. Use smallest size appropriate for 7 day trial.	
	gentamicin mupirocin			
	BENZOYL PEROXIDE/ADAPALENE COMBOS			adapalene/benzoyl peroxide gel 0.1-2.5% (BRAND IS PREFERRED)
	EPIDUO*			
	BENZOYL PEROXIDE/CLINDAMYCIN COMBOS		Clients must be 12 to 20 years of age and have a diagnosis of acne vulgaris. Requires prior authorization for clients less than 12 years of age. Acne combinations are limited to clients under the age of 21.	ACANYA (use preferred agent) clindamycin/benzoyl peroxide 1-5% (BRAND IS PREFERRED) ONEXTON (use preferred agent)
		BENZACLIN* clindamycin/benzoyl peroxide 1.2-5% (Refrig)		
	CORTICOSTEROIDS - STEP 1 AGENTS C=CREAM; G=GEL; L=LOTION; O=OINTMENT		Trial and failure of ALL preferred agents greater than or equal to 14 days in the last 90 days.	PANDEL prednicarbate 0.1% (C,O) TEXACORT 2.5% (S)
	LOW POTENCY			
	alclometasone desonide DESOWEN 0.05% (L) fluocinolone 0.01% hydrocortisone butyrate 0.1% (C) hydrocortisone 1%, 2.5% (C,L,O) SYNALAR 0.01%			
	MEDIUM POTENCY		Trial and failure of ALL preferred agents greater than or equal to 14 days in the last 90 days.	Clocortolone Pivalate CORDRAN/SP fluticasone 0.05% (L) hydrocortisone butyrate 0.1% (O) TOPICORT LP TRIANEX
	betamethasone valerate CUTIVATE 0.05% (C) DERMATOP 0.1% (C) desoximetasone 0.05% (C) ELOCON 0.1% fluocinolone 0.025% fluticasone 0.05% (C) hydrocortisone probutate 0.1% (C) mometasone SYNALAR 0.025% TOPICORT 0.05% (C) triamcinolone 0.025%, 0.1%			

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, **Dosage Limitation List** (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.

THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHARGE HEALTHCARE WITH ANY QUESTIONS</small>
DERMATOLOGY continued	HIGH POTENCY		Trial and failure of ALL preferred agents greater than or equal to 14 days in the last 90 days.	APEXICON amcinonide 0.1% (C,L,O) augmented betamethasone 0.05% (G,L,O) clobetasol 0.05% (L) desoximetasone 0.05%, 0.25% (C,G,O) fluciclonide 0.1% (C) HALOG
	betamethasone dipropionate clobetasol/E 0.05% (C,G,O,S) diflorasone DIPROLENE 0.05% (L) fluciclonide flurandrenolide fluticasone 0.005% (O) halobetasol TEMOVATE/E TOPICORT 0.25% (C) triamcinolone 0.5% ULTRAVATE 0.05%			
	IMMUNOMODULATORS - STEP 2 AGENTS		To receive a step 2 agent: Trial and failure of a preferred medium potency topical corticosteroid greater than or equal to a 21 day trial and a trial and failure of a preferred high potency topical corticosteroid greater than or equal to a 21 day trial in the last 90 days. For clients less than two (2) years of age, a trial and failure of a preferred low potency corticosteroid greater than or equal to a 21 day trial and a trial and failure of a preferred medium potency topical corticosteroid greater than or equal to a 21 day trial in the last 90 days.	
		ELIDEL tacrolimus ointment		
	PHOSPHODIESTERASE 4 INHIBITOR - STEP 3 AGENT		To receive a step 3 agent: Trial and failure of a preferred step 2 agent (immunomodulator) greater than or equal to a 21 day trial within the last 30 days.	DUPIXENT EUCRISA
	PLAQUE PSORIASIS (PP)		Client must have diagnosis of PP prior to approval of a step 1 agent (Enbrel or Humira). To receive Cosentyx, the client must have a diagnosis of PP and a 56-day trial and failure of Humira. To receive Tremfya, the client must have a diagnosis of PP and a 56-day trial and failure of Enbrel. To receive a non-preferred agent, client must have a diagnosis of PP and a 56-day trial and failure of both preferred agents.	OTEZLA REMICADE SILIQ STELARA TALTZ
	STEP 1 AGENTS			
		ENBREL HUMIRA		
	STEP 2 AGENTS			
		COSENTYX TREMFYA		
	SALICYLIC ACID			All other topical salicylic acid formulations.
	salicylic acid cream 6% salicylic acid lotion 6% salicylic acid shampoo 6%			
	SCABICIDES/PEDICULICIDES		Trial and failure of a preferred agent in the last 12 months.	LINDANE OVIDE
NATROBA permethrin SKLICE				
UREA			All other topical urea formulations.	
ALUVEA CREAM 33% UMECTA EMULSION umecta mousse aerosol 40% urea lotion 40% urea lotion 45%				
DIABETES	DIABETES AGENTS			metformin SR 24HR osmotic release (use preferred agent) metformin SR 24HR modified release (use preferred agent) RiOMET (use preferred agent)
	BIGUANIDES			
	metformin/ER			
	α-GLUCOSIDASE INHIBITORS		Trial and failure of metformin and a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	GLYSET*
	acarbose			
	MEGLITINIDES		Trial and failure of metformin and a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	repaglinide
	nateglinide			
	THIAZOLIDINEDIONES		Trial and failure of metformin and a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	ACTOSPUS MET (use separate agents) AVANDIA AVANDAMET (use separate agents)
	pioglitazone			
	SULFONYLUREAS		Trial and failure of metformin and a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	
	glimperide/ER glipizide/ER glyburide/ER			
DIPEPTIDYL PEPTIDASE 4 (DPP-4) INHIBITORS		Trial and failure of metformin greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a preferred agent. A 90 day trial of failure of the preferred agent is required before approval can be give for a non-preferred agent.	alogliptin GLYXAMBI (use separate preferred agents) ONGLYZA TRADJENTA	
	JANUVIA			
DPP-4 INHIBITOR COMBO AGENTS		Trial and failure of metformin greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a preferred agent. A 90 day trial of failure of the preferred agent is required before approval can be give for a non-preferred agent.	alogliptin/metformin alogliptin/pioglitazone (use separate preferred agents) JENTADUETO JUVISYNC (use separate preferred agents) KOMBIGLYZE	
	JANUMET/XR			
INCRETIN MIMETICS (GLP-1 RECEPTOR AGONISTS)		Trial and failure of metformin greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a preferred agent. A 90 day trial of failure of the preferred agent is required before approval can be give for a non-preferred agent.	ADLYXIN BYDUREON SOLIUQA TANZEUM TRULICITY	
	BYETTA VICTOZA			
		Dosage Limits Apply: Victoza: 1.8mg/day		

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria.					
THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHARGE HEALTHCARE WITH ANY QUESTIONS</small>	
DIABETES continued	SGLT2 INHIBITORS		Trial and failure of metformin greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a preferred agent. A 90 day trial of failure of the preferred agent is required before approval can be give for a non-preferred agent.	GLYXAMBI (use separate preferred agents) INVOKAMET/XR(use separate preferred agents) INVOKANA SYNJIARDY/XR (use separate preferred agents) XIGDUO XR (use separate preferred agents)	
	FARXIGA JARDIANCE				
	LONG-ACTING INSULIN		Prior authorization will be required when using two different delivery forms of the same type of insulin concurrently	LANTUS OPTICLIK (use preferred agent) TOUJEO (use preferred agent) TRESIBA (use preferred agent) XULTOPHY (use preferred agent)	
	LANTUS SOLOSTAR LANTUS vial LEVEMIR				
DIABETIC METERS/TEST STRIPS			Quantity limits apply: Insulin Dependent Clients: 10 strips/day Non-Insulin Dependent Clients: 4 strips/day Clients are limited to 1 meter/365 days	ALL OTHER METERS AND TEST STRIPS	
FREESTYLE INSULINX FREESTYLE LITE FREESTYLE FREEDOM LITE ONE TOUCH ULTRA ONE TOUCH ULTRA 2 ONE TOUCH ULTRA MINI ONE TOUCH ULTRASMART ONE TOUCH VERIO ONE TOUCH VERIO FLEX PRECISION XTRA					
FIBROMYALGIA	FIBROMYALGIA STEP 1				
	amitriptyline cyclobenzaprine				
	FIBROMYALGIA STEP 2		Trial and failure of a Step 1 agent greater than or equal to six (6) weeks in the last 12 months is required for approval of a Step 2 agent.		
FIBROMYALGIA STEP 3		Trial and failure of a Step 1 agent and a Step 2 agent greater than or equal to six (6) weeks in the last 12 months is required for approval of a Step 3 agent.			
duloxetine LYRICA			Dosage Limits Apply: Lyrica: 600mg/day		
GASTROINTESTINAL	DIGESTIVE ENZYMES		Prior authorization required.	PANCREAZE pancrelipase PERTZYE TRI-PASE ULTRESA VIKASE	
	CREON ZENPEP				
	PREGNANCY INDUCED NAUSEA/VOMITING				
	DICLEGIS				
	CHRONIC IDIOPATHIC CONSTIPATION		Client must have a diagnosis of chronic idiopathic constipation to receive a preferred agent. To receive a non-preferred agent, the client must have a diagnosis of chronic idiopathic constipation and a 30-day trial and failure of a preferred agent within the last 12 months.	TRULANCE	
	AMITIZA LINZESS				
	IRRITABLE BOWEL SYNDROME WITH CONSTIPATION		Client must have a diagnosis of Irritable Bowel Syndrome (IBS) with constipation.		
	AMITIZA LINZESS				
	OPIOID-INDUCED CONSTIPATION AGENTS		Client must have a diagnosis of opioid-induced constipation and a three (3) month trial and failure of a secretory agent to receive the preferred agent. To receive the non-preferred agent, the client must have a diagnosis of opioid-induced constipation, a three (3) month trial and failure of a secretory agent, and a three (3) month trial and failure of the preferred agent.	MOVANTIK*	
	AMITIZA				
PROTON PUMP INHIBITORS		Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Lansoprazole solutabs will be approved for children less than or equal to 8 years of age.	ACIPHEX SPRINKLES amox/clarith/lanso pack (use separate agents) DEXILANT esomeprazole 24.65mg and 49.3mg NEXIUM* omeprazole 20.6mg capsules (use preferred agent) omeprazole tablets (use preferred agent) omeprazole/sodium bicarbonate OMECLAMOX (use separate agents) PREVACID solutabs rabeprazole VIMOVO (use separate agents)		
lansoprazole capsules omeprazole capsules pantoprazole					
MESALAMINE		Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	APRISO ASACOL HD CANASA DELZICOL GIAZO mesalamine DR tab 1.2gm (BRAND IS PREFERRED) SFROWASA		
LIALDA* mesalamine enema PENTASA					
GOUT	COLCHICINE			COLCRYS (use preferred agent) MITIGARE (use preferred agent)	
	colchicine				
XANTHINE OXIDASE AND URAT1 INHIBITORS		Trial and failure of a preferred agent greater than or equal to a 60 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	ZURAMPIC*		
allopurinol ULORIC			*Concurrent use of a preferred agent will be required with Zurampic.		

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, **Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.**

Therapeutic Class	Preferred Agents	Preferred Agents Requiring Clinical Criteria	Clinical Criteria	Non-Preferred Agents Generic Mandatory Policy Applies <small>PLEASE CONTACT YOUR HEALTHCARE PROVIDER WITH ANY QUESTIONS</small>
HEMATOLOGY	LOW MOLECULAR WEIGHT HEPARIN (LMWH)		Prior authorization will be required for the 300mg/3ml strength.	FRAGMIN (<i>use preferred agent</i>) LOVENOX 300MG/3ML*
	enoxaparin			
	DIRECT THROMBIN INHIBITOR		Client must have diagnosis of non-valvular atrial fibrillation and relative contraindication to warfarin for approval, treatment for deep vein thrombosis (DVT) or pulmonary embolism (PE), or for the reduction in the risk of recurrence of DVT and PE after initial therapy.	
		PRADAXA		
	FACTOR XA INHIBITOR		Limited to being used for the prophylaxis of venous thromboembolism (VTE) in adult patients hospitalized for an acute medical illness who are at risk for thromboembolic complications due to moderate or severe restricted mobility and other risk factors for VTE	
		BEVYXXA		
	SELECTIVE FACTOR XA INHIBITOR		Client must have diagnosis of non-valvular atrial fibrillation, treatment for deep vein thrombosis (DVT) prophylaxis in knee or hip replacement, treatment of DVT and pulmonary embolism (PE), and for the reduction in the risk of recurrent DVT and PE after initial therapy.	SAVAYSA (<i>use preferred agent</i>)
		ELIQUIS XARELTO		
	THIENOPYRIDINE DERIVATIVES		Prior authorization required for clients on antiplatelet therapy greater than one (1) year.	
	clopidogrel EFFIENT ticlopidine			
	CPTP DERIVATIVES		Prior authorization is required.	BRILINTA
PAR-1 ANTAGONIST		Client must have diagnosis of reduction of thrombotic cardiovascular events with a history of myocardial infarction (MI) or with peripheral arterial disease (PAD). Must be used in conjunction with aspirin or clopidogrel.		
	ZONTIVITY			
ANTIHEMOPHILIC FACTOR VIII			NOVOEIGHT (<i>requires prior authorization</i>)	
ADVATE ADYNOVATE AFSTYLA ELOCTATE HELIXATE FS HEMOPIL M KOATE/KOATE-DVI KOGENATE FS/BIO-SET KOVALTRY MONOCLATE-P NUWIQ OBIZUR RECOMBINATE XYNTHA/SOLOFUSE				
ANTIHEMOPHILIC FACTOR/VWF				
ALPHANATE HUMATE-P WILATE				
HEPATITIS C	DIRECT ACTING ANTIVIRALS		Limited to FDA approved indication. Prior authorization will be required prior to use of preferred agents. **Positive SVR 12 will be required for consideration for retreatment Please submit PA requests on the Hepatitis C PA form available at www.wymedicaid.org .	DAKLINZA (<i>use preferred agent</i>) OLYSIO (<i>use preferred agent</i>) SOVALDI (<i>use preferred agent</i>) TECHNIVIE (<i>use preferred agent</i>) VIEKIRA PAK/XR (<i>use preferred agent</i>) ZEPATIER (<i>use preferred agent</i>)
		EPCLUSA HARVONI MAVYRET** VOSEVI**		
HIDRADENITIS SUPPURATIVA	IMMUNOMODULATORS		Humira will not be covered as a first line agent for the diagnosis of hidradenitis suppurativa.	
		HUMIRA		
HORMONES	GROWTH HORMONE		PA is required for use outside of FDA-approved indications. Evaluation by an endocrinologist is preferred. Clinical evidence of improved growth will be required on a yearly basis to support ongoing utilization. Clinical evidence of need for growth hormone will be required for adult growth hormone deficiency and pediatric growth failure due to inadequate endogenous growth hormone. Trial and failure of two (2) preferred agents within the last 12 months will be required for the following indications: Pediatric: Growth failure due to inadequate endogenous growth hormone, Prader-Willi syndrome, children born small for gestation, Turner syndrome. Adult: Replacement for those with growth hormone deficiency.	HUMATROPE OMNITROPE SAIZEN SEROSTIM TEV-TROPIN ZORBIVE
		GENOTROPIN NORDITROPIN NUTROPIN AQ		
	PROGESTIN		Prior authorization is required.	
		MAKENA		
	TESTOSTERONE TOPICAL GELS		Testosterone agents are only allowed for diagnosis of hypogonadism or insufficient testosterone production. <i>Other testosterone dosage form products will require a diagnosis of hypogonadism or insufficient testosterone production (not outlined on PDL).</i>	NATESTO NASAL GEL (<i>use preferred agent</i>) TESTIM GEL (<i>use preferred agent</i>) testosterone gel 1% (BRAND IS PREFERRED) testosterone gel 2% (<i>use preferred agent</i>) VOGELXO GEL (<i>use preferred agent</i>)
		ANDROGEL*		

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, **Dosage Limitation List** (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.

THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHANGING HEALTHCARE WITH ANY QUESTIONS</small>
HORMONES continued	<p style="text-align: center;">ORAL CONTRACEPTIVES</p> altavera alyacen 1-35, 7/7/7 amethyst azurette apri aubra aviane balziva bekyree blisovi 1-20 FE/24, 1.5-30 FE briellyn camila caziant chateal cyclofem 1-35, 7/7/7 cyred cryselle dasetta 1-35, 7/7/7 debilitane delyla DESOGEN deso/ethinyl estradiol elinest emoquette enpresse enskyce errin estarvila falmina FEMCON FE CHEWABLE gianvi gildagia gildess 1-20/FE/24, 1.5-30/FE heather jencycla jolessa jolivet juleber junel 1-20/FE/24, 1.5-30/FE kariva kelnor kimidess kurvelo larin 1-20/FE/24, 1.5-30/FE leena lessina levonest levonor/ethi levora lomedra 24 FE LOSEASONIQUE* low-ogestrel lutera lyza marlissa microgestin 1-20/FE/24, 1.5-30/FE MODICON mono-linyah mononessa myzila NECON 0.5-35, 1-35, 7/7/7, 10/11-28 nora-be norgest/ethinyl estradiol norethindrone norlyroc noreth/ethin 1-20/FE/24 NORINYL 1/50-28 nortrel 0.5-35, 1-35, 7/7/7 ocella OGESTREL orsythia ORTHO TRI-CYCLEN LO* ORTHO-NOVUM 1/35, 7/7/7* phillith pimtrea pirmella 1-35, 7/7/7 portia previfem reclippen SEASONIQUE* setlakin sprintec sharobel sronyx syeda tilia FE tri-estaryl tri-legest FE tri-linyah trinessa TRI-NORINYL* tri-previfem tri-sprintec trivora velivet vestura vienna viorele vyfemla wera 0.5-35 YAZ zarah zenchent ZOVIA			amethia/LO (BRAND IS PREFERRED) aranelle (use preferred agent) ashlyna (BRAND IS PREFERRED) BEYAZ (PA required) BREVICON (use preferred agent) camrese/LO (BRAND IS PREFERRED) daysee (BRAND IS PREFERRED) drospir/ethi (use preferred agent) estarylla tri-lo (BRAND IS PREFERRED) FALESSA KIT (use preferred agent) introvale (use preferred agent) layolis FE chewable (PA required) levonorgest/ethinyl estrad (91-Day) levonorgest/ethinyl estradiol (Continuous) (use preferred agent) levonorgest/ethinyl estradiol/LO (84-7) (BRAND IS PREFERRED) LO LOESTRIN (PA required) LO MINASTRIN FE (PA required) loryna (use preferred agent) MINASTRIN 24 FE CHEWABLE (PA required) NATAZIA (PA required) norgest/ethi estradiol lo (BRAND IS PREFERRED) NATAZIA (PA required) NECON 1/50-28 (use preferred agent) nikki (use preferred agent) noreth/ethin FE chewable (PA required) NORINYL 1/35 (use preferred agent) quasense (use preferred agent) QUARTETTE (PA required) SAFYRAL (PA required) tri-lo sprintec (BRAND IS PREFERRED) trinessa lo (BRAND IS PREFERRED) wymzya FE chewable (BRAND IS PREFERRED) zenchent FE chewable (BRAND IS PREFERRED)

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.

THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHARGE HEALTHCARE WITH ANY QUESTIONS</small>	
HYPERLIPIDEMIA	BILE ACID SEQUESTERANT		Trial and failure of ALL preferred agents greater than or equal to six (6) months in the last 12 months will be required before approval can be given for a non-preferred agent.	WELCHOL	
	cholestyramine/light colestipol				
	STATINS, LOW POTENCY			Trial and failure of a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. If client's current medication therapy is contraindicated with the preferred statin(s) due to a drug-drug interaction, a non-preferred agent may be obtained with a prior authorization. Prior authorization will be required for clients under the age of 10.	fluvastatin/ER
	lovastatin pravastatin				
	STATINS, HIGH POTENCY			Trial and failure of a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. If client's current medication therapy is contraindicated with the preferred statin(s) due to a drug-drug interaction, a non-preferred agent may be obtained with a prior authorization. Prior authorization will be required for clients under the age of 10.	LIVALO rosuvastatin
	atorvastatin simvastatin				
STATIN COMBINATIONS		Trial and failure of a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Prior authorization will be required for clients under the age of 10.	amlodopine/atorvastatin (BRAND IS PREFERRED) ezetimibe-simvastatin (BRAND IS PREFERRED)		
CADUET* VYTORIN*					
TRIGLYCERIDE LOWERING AGENTS		Trial and failure of a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	ANTARA fenofibric fenofibrate 43, 50, 120, 130, and 150me LIOPEN omega-3-acid VASCEPA		
fenofibrate 48, 54, 67, 134, 145, 160, and 200mg gemfibrozil					
HYPERTENSION	ANGIOTENSIN RECEPTOR BLOCKERS (ARBs)		Non-preferred ARBs will require a history of ALL preferred ARBs before approval can be given.	candesartan eprosartan 600mg TEVETEN 400mg	
	EDARBI irbesartan losartan olmesartan telmisartan valsartan				
	ARBs AND DIURETICS			Non-preferred ARB/diuretic combinations will require a history of ALL preferred ARBs before approval can be given.	candesartan HCTZ telmisartan HCTZ TEVETEN HCTZ
	EDARBYCLOR irbesartan HCTZ losartan HCT olmesartan HCTZ valsartan HCTZ				
ALPHA-BLOCKERS			clonidine patch (BRAND IS PREFERRED) NEXICLON XR (use preferred agent)		
CATAPRES PATCHES* clonidine					
INFECTIOUS DISEASE	QUINOLONES		*Tobi Podhaler requires a 28 day trial of a preferred agent, as well as 28 days off of that same preferred agent prior to approval. Minimum day supply of at 56 days is required	FACTIVE moxifloxacin NOROXIN PROQUIN	
	ciprofloxacin/ER levofloxacin ofloxacin				
	DOXYCYCLINE			ADOXA (use preferred agent) DORYX (use preferred agent) ORACEA (use preferred agent)	
	doxycycline				
	MINOCYCLINE			SOLODYN (use preferred agent)	
	minocycline/ER				
	INHALED TOBRAMYCIN			inhaled tobramycin (use preferred agent)	
	BETHKIS KITABIS	TOBI PODHALER*			
KEFLEX		cephalexin 750mg (BRAND IS PREFERRED)			
KEFLEX 750mg*					
ANTI-RETROVIRALS					
DESCOVY EVOTAZ GENVOYA NORVIR ODEFSEY PREZCOBIX					

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, **Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.**

Therapeutic Class	Preferred Agents	Preferred Agents Requiring Clinical Criteria	Clinical Criteria	Non-Preferred Agents Generic Mandatory Policy Applies <small>PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small>
INFLAMMATION	NSAIDs		Trial and failure of two (2) preferred agents each greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Dosing and quantity limits apply for ketorolac (limit 5days/34 days; max dose 40mg/day for oral tablets).	CALDOLOR (use preferred agent) CAMBIA POWDER (use preferred agent) celecoxib diclofenac 1.5% solution (additional criteria applies) diclofenac 3% gel (additional criteria applies) fenoprofen FLECTOR (additional criteria applies) mefenamic acid NEOPROFEN (use preferred agent) SPRIX (additional criteria applies) TIVORBEX (use preferred agent) VIVLODEX (use preferred agent) VOLTAREN* (additional criteria applies) ZIPSOR (use preferred agent) ZORVOLEX (use preferred agent)
	diclofenac tablets etodolac flurbiprofen ibuprofen indomethacin ketoprofen ketorolac meclufenamate meloxicam nabumetone naproxen oxaprozin piroxicam sulindac tolmetin			
	ORAL CORTICOSTEROIDS			CELESTONE (use preferred agent)
	budesonide cortisone acetate dexamethasone/intensol hydrocortisone methylprednisone prednisolone prednisone			
INSOMNIA	NON-BENZODIAZEPINES		Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Prior authorization will be required for clients under the age of 18. Rozerem is non-preferred without a history of substance abuse Prior authorization will be required when a client is taking more than one insomnia agent concurrently. Dosage limits apply: zaleplon: 30mg/day zolpidem: 15mg/day	BELSOMRA EDLUAR (additional criteria applies) eszopiclone INTERMEZZO (additional criteria applies) ROZEREM zolpidem ER ZOLPIMIST (additional criteria applies)
	zaleplon zolpidem			
MENTAL HEALTH	ALZHEIMER AGENTS		Client must have a diagnosis of dementia. Trial and failure of two (2) preferred agents greater than or equal to six (6) weeks WITHIN THE LAST 2 YEARS will be required before approval can be given for a non-preferred agent. One of the trials of preferred agents must be in the same class (NaSS, NDRI, SSRI, or SNRI) as the requested non-preferred agent.	donepezil/ODT EXELON PATCH* galantamine/ER memantine tablets/solution rivastigmine capsules mirtazapine 7.5mg and rapid dissolve tablets (use preferred agent) APLENZIN FORFIVO XL fluoxetine tablets (use preferred agent) VIIBRYD duloxetine** desvenlafaxine FETZIMA venlafaxine ER tablets (use preferred agent) TRINTELLIX***
	ANTIDEPRESSANTS			
	NORADRENERGIC/SPECIFIC SEROTONERGICS (NaSS)			
	mirtazapine 15, 30, and 45mg			
	NOREPINEPHRINE/DOPAMINE REUPTAKE INHIBITORS (NDRI)			
	bupropion ER/SR/XL			
	SELECTIVE SEROTONIN REUPTAKE INHIBITORS (SSRI)			
	citalopram escitalopram fluoxetine capsules paroxetine IR/CR sertraline			
	SEROTONIN/NORPINEPHRINE REUPTAKE INHIBITORS (SNRI)			
	venlafaxine ER capsules			

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, **Dosage Limitation List** (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.

THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small>
MENTAL HEALTH continued	ATYPICAL ANTIPSYCHOTICS ABILIFY MAINTENA ABILIFY ODT* aripiprazole tab/solution ARISTADA FANAPT INVEGA* INVEGA SUSTENNA/TRINZ LATUDA*** olanzapine quetiapine RISPERDAL CONSTA risperidone SAPHRIS ziprasidone ZYPREXA RELPREVV		**Quetiapine doses less than 100mg will require prior authorization without a diagnosis of mood disorder or major depressive disorder. For titration doses, contact the Change Healthcare Pharmacy Help Desk for an override. Clients five (5) years of age and younger will require prior authorization before approval. **Trial and failure of two (2) preferred agents greater than or equal to 30 days in the last 12 months will be required before approval can be given for REXULTI or VRAYLAR. ***Clients twelve (12) years of age and younger will require a prior authorization to receive approval of Latuda. Dosage limits apply: -aripiprazole <13 years of age: 23mg/day -aripiprazole ≥13 years of age: 45mg/day -FANAPT: 36mg/day -INVEGA: 18mg/day -LATUDA 13-17 years of age: 120mg/day -LATUDA >17 years of age: 240mg/day -olanzapine <13 years of age: 15mg/day -olanzapine ≥13 years of age: 30mg/day -quetiapine <13 years of age: 600mg/day -quetiapine 13-17 years of age: 900mg/day -quetiapine >17 years of age: 1200mg/day -risperidone ≤ 17 years of age: 5mg/day -risperidone >17 years of age: 24mg/day -SAPHRIS: 30mg/day -ziprasidone ≤17 years of age: 180mg/day -ziprasidone >17 years of age: 300mg/day	aripiprazole ODT (BRAND IS PREFERRED) paliperidone (BRAND IS PREFERRED) REXULTI* quetiapine XR (use preferred agent) VRAYLAR**
	SPECIAL ATYPICAL ANTIPSYCHOTICS		Dosage limits apply: 1350mg/day	VERSACLOZ Suspension (use preferred agent)
	clozapine/ODT	AMPHETAMINES LONG ACTING AMPHETAMINES amphetamine salts combo XR dextroamphetamine CR caps VYVANSE CAPSULES**		Clients over the age of 17 must have a diagnosis of ADD, ADHD, narcolepsy, obstructive sleep apnea, shift work sleep disturbance, MS fatigue (see MS Fatigue criteria below), or refractory depression (see refractory depression criteria below).
IMMEDIATE RELEASE AMPHETAMINES		amphetamine salts combo dextroamphetamine tablets	Diagnosis of MS fatigue will require a fatigue severity scale score of 5.0, a 60-day trial of amantadine and discontinuation of medications that may contribute to drowsiness and fatigue.	
METHYLPHENIDATES LONG ACTING METHYLPHENIDATES		DAYTRANA FOCALIN XR* methylin ER methylphenidate ER/CR/SA/SR tablets***	Diagnosis of refractory depression will require a 6-week trial and failure of an antidepressant (monotherapy) and continued concomitant use of an antidepressant with the stimulant.	METHYLPHENIDATES APTENSIO XR COTEMPLA dexmethylphenidate ER (BRAND IS PREFERRED) methylphenidate ER/CR/SR capsules (METADATE CD/RITALIN LA) QUILLICHEW QUILLIVANT XR SUSPENSION
IMMEDIATE RELEASE METHYLPHENIDATES		dexmethylphenidate methylin tablets methylphenidate tablets	Prior Authorization will be required for clients under the age of 4. **Vyvanse will be approved for the diagnosis of binge-eating disorder for clients 18 years of age and older. Authorizations will be approved for 12 weeks, and further use of Vyvanse for this diagnosis will require additional documentation prior to approval. Claims will require Prior Authorization if clients have a history of the following: glaucoma, cardiac arrhythmias, arteriosclerosis, untreated hypertension, untreated hyperthyroidism, substance abuse, or current MAO inhibitor use. Trial and failure of two (2) preferred agents (each from a different class: methylphenidate and amphetamine) greater than or equal to a 30 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Dosage limits apply: amphetamine salts combo XR: 60mg/day amphetamine salts combo: 60mg/day amphetamine salts combo (narcolepsy): 90mg/day DAYTRANA: 45mg/9 hour patch/day dextroamphetamine: 90mg/day dextroamphetamine CR: 90mg/day dexmethylphenidate: 30mg/day FOCALIN XR < 13 years of age: 45mg/day FOCALIN XR > 13 years of age: 60mg/day methylin/methylphenidate/ER: 90mg/day VYVANSE: 105mg/day	

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.

Therapeutic Class	Preferred Agents	Preferred Agents Requiring Clinical Criteria	Clinical Criteria	Non-Preferred Agents Generic Mandatory Policy Applies <small>PLEASE CONTACT CHARGE HEALTHCARE WITH ANY QUESTIONS</small>
MENTAL HEALTH continued	SELECTIVE ALPHA-ADRENERGIC AGONIST		To obtain the non-preferred agent , client must meet the following criteria: Client must have a diagnosis of ADD or ADHD Prior authorization will be required for clients under the age of 4. To receive Kapvay, clients must have completed a 14 day trial of clonidine IR with benefit in the previous 12 months.	KAPVAY*
	clonidine			
	GUANFACINE AGENTS		To obtain the non-preferred agent , client must meet the following criteria: Client must have a diagnosis of ADD or ADHD Prior authorization will be required for clients under the age of 4. To receive guanfacine ER, clients in the previous 12 months must have: A) a trial and failure of a stimulant greater than or equal to a 14 day supply, or B) a trial and failure of Strattera greater than or equal to a 30 day supply, or C) a contraindication to ADHD medications (including stimulant and non-stimulant), or D) a diagnosis of a TIC disorder, AND E) a 14 day trial of guanfacine with benefit	guanfacine ER
guanfacine				
	SELECTIVE NOREPINEPHRINE REUPTAKE INHIBITOR		Clients must have a diagnosis for ADD, ADHD, narcolepsy, obstructive sleep apnea, shift work sleep disturbance, or refractory depression (see refractory depression criteria below). Diagnosis of refractory depression will require a 6-week trial and failure of an antidepressant (monotherapy) and continued concomitant use of an antidepressant with the stimulant. Prior Authorization will be required for clients under the age of 4. Claims will require Prior Authorization if clients have a history of the following: glaucoma, cardiac arrhythmias, arteriosclerosis, untreated hypertension, untreated hyperthyroidism, substance abuse, or current MAO inhibitor use. Dosage limits apply: atomoxetine: 150mg/day	
		atomoxetine		
MIGRAINE	TRIPTANS		Trial and failure of two preferred agents will be required for approval of a non-preferred agent. Rizatriptan will be approved for clients between 6 and 17 years of age Quantity limits apply: naratriptan 1mg: 25 tabs/34 days naratriptan 2.5mg: 10 tabs/34 days RELPAK 20mg: 20 tabs/34 days RELPAK 40mg: 14 tabs/34 days sumatriptan vials: 2 vials/34 days sumatriptan nasal: 6 bottles/34 days sumatriptan 25mg: 41 tabs/34 days sumatriptan 50mg: 20 tabs/34 days sumatriptan 100mg: 10 tabs/34 days	almotriptan frovatriptan ONZETRA (use preferred agent) rizatriptan TREMIMET ZEMBRACE (use preferred agent) zolmitriptan
	naratriptan RELPAK sumatriptan			
MULTIPLE SCLEROSIS	STEP 1 MS AGENTS		Trial and failure of one injectable preferred agent will be required before approval can be given for the step 2 MS agent (Gilenya).	AUBAGIO COPAXONE 40MG/ML (use preferred agent) EXTAVIA LEMTRADA OCREVUS* PLEGRIDY TECHIDERA TYSABRI (additional criteria applies) ZINBRYTA
	IMMUNOMODULATOR (GLATIRAMER INJECTION)			
	COPAXONE 20MG/ML			
	INTERFERON			
	AVONEX BETASERON REBIF		Trial and failure of a two preferred agents (each from a separate class) will be required before approval can be given for a non-preferred agent. *Ocrevus will be approved for a diagnosis of primary progressive multiple sclerosis. For relapsing forms of multiple sclerosis, the requirements listed above will need to be followed	
	STEP 2 MS AGENTS		For Tysabri, in addition to the above criteria, additional prior authorization criteria applies.	
		GILENYA		
NEUROPATHIC PAIN	TRICYCLIC ANTIDEPRESSANTS		For the diagnosis of neuropathic pain, trial and failure of a tricyclic antidepressant greater than or equal to a 12 week supply AND trial and failure of gabapentin at a dose of 3600mg per day for greater than or equal to a 12 week supply in the last 12 months will be required before approval can be given for a non-preferred agent.	duloxetine LYRICA
		amitriptyline desipramine imipramine nortriptyline		
	GABAPENTIN			
		gabapentin		

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.

THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHARGE HEALTHCARE WITH ANY QUESTIONS</small>
OPHTHALMICS	OP. -ANTI-ALLERGICS		Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Emadine, Alomide, and Alocril will be approved for pregnancy. Alomide will be approved for children under the age of 3.	ALAMAST ALOCRIL ALOMIDE ALREX azelastine BEPREVE EMADINE epinastine ketotifen LASTACAFT olopatadine 0.1% and 0.2%
	OP. -ANTIBIOTICS- QUINOLONES		Trial and failure of a preferred agent greater than or equal to 5 days in the last 12 months will be required before approval can be given for a non-preferred agent. Azasite will be approved for pregnancy.	AZASITE BESIVANCE gatifloxacin IQUIX levofloxacin moxifloxacin 0.5% (BRAND IS PREFERRED) ZYMAR
	OP. -ANTI-INFLAMMATORY		Trial and failure of ALL preferred agents each greater than or equal to 5 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.	ACUJAR/LS/PF (use preferred) ACUVAIL bromfenac 0.9% BROMSITE DUREZOL NEVENAC PROLENSA
	OP. -BETA-BLOCKERS		Trial and failure of three (3) preferred agents each greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Betoptic S will be approved for those with heart and lung conditions.	BETIMOL BETOPTIC S ISTALOL
	OP. -CARBONIC ANHYDRASE INHIBITOR		Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent.	AZOPT
	OP. - COMBO PRODUCTS		Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent.	
	OP. - DRY EYE AGENTS		Trial and failure of the preferred agent greater than or equal to 12 weeks will be required before approval can be given for the non-preferred agent.	RESTASIS MULTIDOSE (use preferred) XIIDRA
	OP. -PROSTAGLANDINS		Trial and failure of ALL preferred agents each greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent.	bimatoprost LUMIGAN 0.1% ZIOPTAN
	OP. -SYMPATHOMIMETICS		Trial of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent.	brimonidine 0.15% (BRAND IS PREFERRED)
	OP. -ALPHAGAN P 0.15%*			
OSTEOPOROSIS	BISPHOSPHONATES		Trial and failure of a preferred agent greater than or equal to 12 months will be required before approval can be given for a non-preferred agent. Fosamax liquid will be approved for clients that have difficulty swallowing.	risedronate ATELVIA FOSAMAX-D ibandronate TYMLOS
	NASAL CALCITONIN			
OTIC	CIPRODEX Neo/Poly/HC Suspension and Solution	ANTIBIOTIC/STEROID COMBINATION		ciprofloxacin 0.2% (use preferred agent) CIPRO HC (use preferred agent) COLY-MYCIN S (use preferred agent) CORTISPORIN-TC (use preferred agent) FLUCINOLONE ACET OIL 0.01% (use preferred agent) ofloxacin (use preferred agent)
OVERACTIVE BLADDER		OVERACTIVE BLADDER AGENTS		Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Oxytrol will be approved for clients that have an inability to swallow.

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, **Dosage Limitation List** (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.

THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHARGE HEALTHCARE WITH ANY QUESTIONS</small>	
PAIN	morphine sulfate ER <u>tablets</u>	<p align="center">LONG-ACTING C-III's</p> <p>fentanyl patch 12.5, 25, 50, 75, and 100mg</p>	<p>Trial and failure of a preferred agent(s) greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.</p> <p>C-III's and C-IV's that are not included on the PDL and are available without prior authorization with the exception of Butrans (generic substitution is mandatory).</p> <p>Concurrent use of a narcotic and benzodiazepine will require prior authorization</p> <p>Fentanyl patches will require a prior authorization unless a client has a cancer diagnosis or previous treatment of at least a 10 day supply within the last 45 days</p> <p>**Butrans requires a trial of morphine sulfate ER or low dose trial of fentanyl patch.</p> <p>***Nucynta ER will be allowed for diabetic peripheral neuropathy or clients with significant gastrointestinal concerns with other CII narcotics.</p> <p>****In addition to above criteria, Embeda requires a diagnosis of drug/substance abuse.</p> <p>Belbuca: 1.2mg/day (1200mcg/day) Butrans: 20mcg, 1 strength at a time, 1 patch every 7 days Fentanyl: 50mcg, 1 strength at a time, 1 patch every 3 days Hysingla ER: 120mg/day Hydromorphone ER: 30mg/day Morphabond: 120mg/day Morphine ER: 120mg/day Methadone: Limited to 3 tablets per day Nucynta ER: 327mg/day Oxycontin: 80mg/day Oxymorphone ER: 40mg/day Xartemis XR: 80mg/day Xtampza ER: 80mg/day Zohydro ER: 120mg/day</p> <p>Clients will be limited to one long-acting narcotic at a time</p>	<p>AVINZA BELBUCA BUTRANS** EMBEDA**** fentanyl patch 37.5, 62.5, 87.5mg hydromorphone ER HYSINGLA ER (additional criteria applies) KADIAN 200mg (use preferred agent) METHADONE MORPHABOND morphine sulfate ER capsules (use preferred) NUCYNTA ER*** oxymorphone ER OXYCONTIN XARTEMIS XR (additional criteria applies) XTAMPZA ER (additional criteria applies) ZOHYDRO ER (additional criteria applies)</p>	
		<p align="center">SHORT-ACTING C-III's</p> <p>codeine sulfate hydrocodone/APAP hydrocodone/IBU hydromorphone LORTAB ELIXIR 10-300MG meperidine morphine sulfate oxycodone oxycodone/APAP oxycodone/ASA</p>		<p>Trial and failure of three (3) preferred agents greater than or equal to a 6 day supply in the last 90 days will be required before approval can be given for a non-preferred agent.</p> <p>*Nucynta will be allowed for diabetic peripheral neuropathy or clients with significant gastrointestinal concerns with other CII narcotics.</p> <p>Concurrent use of a narcotic and benzodiazepine will require prior authorization</p> <p>All short-acting narcotics, after 42 days of consecutive use of any combination of short-acting narcotics, will be limited to 4 tablets per day (liquids have specific dosing limits per medication - please refer to dosage limitation chart at www.wymedicaid.org)</p> <p>Clients will be limited to one short-acting narcotic at a time</p>	<p>levorphanol NUCYNTA* oxymorphone oxycodone/IBU</p>
		<p align="center">C-III/C-V AGENTS</p> <p>tramadol</p>		<p>Trial and failure of a preferred agent(s) greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.</p> <p>Quantity and dosage limits apply (max 8 tabs/day).</p> <p>**Butrans will require a 14 day trial and failure of tramadol IR and a 14 day trial and failure of tramadol ER prior to approval</p>	<p>BUTRANS** RYBIX ODT tramadol/apap tramadol ER capsules tramadol ER tablets</p>
PHOSPHATE BINDERS	<p align="center">PHOSPHATE BINDERS</p> <p>calcium acetate RENAGEL</p>		<p>Prior authorization required for non-preferred agents.</p>	<p>AURYXIA lanthanum PHOSLYRA sevelamer VELPHORO</p>	
PROSTATE	<p align="center">5-ALPHA-REDUCTASE INHIBITORS</p> <p>finasteride</p>		<p>Trial and failure of a preferred agent greater than or equal to a 30 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.</p>	<p>dutasteride dutasteride/tamsulosin (use separate agents)</p>	
	<p align="center">ALPHA BLOCKERS</p> <p>doxazosin tamsulosin terazosin</p>		<p>Trial and failure of a preferred agent greater than or equal to a 30 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.</p>	<p>alfuzosin dutasteride/tamsulosin (use separate agents) RAPAFLO</p>	

WYOMING MEDICAID
Preferred Drug List (PDL) - January 24, 2018

Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria.					
THERAPEUTIC CLASS	PREFERRED AGENTS	PREFERRED AGENTS REQUIRING CLINICAL CRITERIA	CLINICAL CRITERIA	NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>PLEASE CONTACT CHANGING HEALTHCARE WITH ANY QUESTIONS</small>	
PULMONARY ANTIHYPERTENSIVES	5-ALPHA-REDUCTASE INHIBITORS		Prior authorization required. Client must have a diagnosis of pulmonary hypertension with documented right-heart catheterization validating the diagnosis.		
		ADCIRCA REVATIO SUSPENSION sildenafil (Revatio A/B rated generic)			
	ENDOTHELIN RECEPTOR ANTAGONISTS		Prior authorization required. Client must have a diagnosis of pulmonary hypertension with documented right-heart catheterization validating the diagnosis.		OPSUMIT (use preferred agent)
		LETAIRIS TRACLEER			
	PROSTACYCLINE VASODILATORS		Prior authorization required. Client must have a diagnosis of pulmonary hypertension with documented right-heart catheterization validating the diagnosis.		
		ORENITRAM			
			Prior authorization required.	UPTRAVI (use preferred pulmonary HTN agent)	
RESTLESS LEG SYNDROME	RESTLESS LEG SYNDROME		Client must have a diagnosis of Restless Leg Syndrome (RLS). Trial and failure of gabapentin greater than or equal to 60 days and a trial and failure of a dopamine agonist greater than or equal to 60 days in the last 12 months will be required before approval can be given for a non-preferred agent. *Neupro will be approved for clients with difficulty swallowing or for clients with a diagnosis of Parkinson's Disease.	HORIZANT NEUPRO*	
		gabapentin pramipexole ropinirole			
SKELETAL MUSCLE RELAXANTS	MUSCLE RELAXANTS		Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months, along with a medical diagnosis of muscle spasticity will be required before approval can be given for a non-preferred agent. Cyclobenzaprine will require a prior authorization for clients concurrently taking a tricyclic antidepressant.	carisoprodol chlorzoxazone cyclobenzaprine ER metaxalone methocarbamol orphenadrine tizanidine capsules (use preferred agent) Carisoprodol is limited to 84 tabs/365 days	
	baclofen cyclobenzaprine tizanidine tablets				
ULCERATIVE COLITIS	IMMUNOMODULATORS		Client must have diagnosis of UC prior to approval of a preferred agent. To receive a non-preferred agent, client must have a diagnosis of UC and a 56-day trial and failure of the preferred agent.	REMICADE	
UVEITIS	IMMUNOMODULATORS		Client must have diagnosis of non-infectious intermediate, posterior, and panuveitis in adult patients		
		HUMIRA			